



# JURNAL MIDPRO

ISSN : 2086-2792 (Print) ISSN : 2684-6764 (Online)

Sekolah Vokasi | Program Studi Diploma III Kebidanan | Universitas Islam Lamongan



- HOME
- ABOUT
- LOGIN
- REGISTER
- SEARCH
- CURRENT
- ARCHIVES
- INDEXING
- ANNOUNCEMENT
- JOURNAL HISTORY

Home > Archives > Vol 12, No 2 (2020)

## Vol 12, No 2 (2020)

JURNAL MIDPRO

DOI: <https://doi.org/10.30736/md.v12i2>

### Table of Contents

Qualitative and Quantitative Analysis of Hydroquinone in Face Cream with TLC and HPLC Methods <i>Rahma Dona, Selvia Selvia</i>	PDF 157-162
Implementation of Online Learning During the Covid-19 Pandemic Period in Supporting the Physical Distancing Policy at Stikes Rajekwesi Bojonegoro in 2020 <i>Nur Azizah, Wiwik Muhidayati, Erien Luthfia, Rina Utami</i>	PDF 163-169
The Effect Of Brain Gym With The Level Of Stress On Elders In Panti Sosial Lanjut Usia Potroyudan Jepara <i>Galia Wardha Alvita, Sholihul Huda</i>	PDF 170-176

#### OPEN JOURNAL SYSTEMS

- Editorial Board
- Reviewers
- Peer Review Process
- Section Policies
- Focus and Scope
- Author Guidelines
- Publication Ethics
- Online Submission
- Open Access Policy
- Author(s) fee
- Copyright Notice
- Login
- Contact



 <i>Galia Wardha Alvita, Sholihul Huda</i>	
<b>Healthy Living To Dismiss Covid-19 In Economic Crisis</b>  <i>R Mochamad A</i>	PDF 177-190
<b>Relationship between Training Intensity and Doctor's Job Satisfaction at Primary Health Care in Indonesian</b>  <i>Gita Purnamasari, Misnaniarti Misnaniarti</i>	PDF 191-197
<b>Phytochemical Screening And Tyrosinase Inhibition Activity Of Leaves Cassia siamea L.</b>  <i>Munawarothus Sholikha, Ainun Wulandari</i>	PDF 198-205
<b>Relationship of Body Mass Index and Haemoglobin with Cardiorespiration Endurane in Female College Students</b>  <i>Nadia Ayu Kusuma Astuti, Faridah Faridah, Iis Fatimawati, Yoga Kertapati</i>	PDF 206-211
<b>Trancendental Meditation Control Blood Sugar Levels On Elderly People</b>  <i>Santi Damayanti, I Kadek Indra Sucipta, Rizky Erwanto</i>	PDF 212-218
<b>Inpatient Satisfaction To Attribute of Health Service Quality Dimensions in Hospitals in Indonesian</b>  <i>Sri Agustini, Haerawati Idris, Misnaniarti Misnaniarti</i>	PDF 219-232
<b>Antenatal Clinical Hypnosis and Birth Ball Exercise as Adjunct Therapy For Management of Labor and Delivery Pain: a Summary of Its Empirical Findings</b>  <i>Ni Gusti Ayu Pramita Aswitami, Ni Gusti Ayu Pitria Septiani</i>	PDF 233-245
<b>Serum Caspase-3 And Tissue Of Cervical Intraepithelial Neoplasia (CIN) In Wistar Rats With Diethylstilbestrol Induction</b>  <i>Sri Rahayu, Iting Iting</i>	PDF 246-252
<b>Influence Akupresur At The Li 4 And Sp 6 To A Decrease In Pain Dysminorrhea</b>  <i>Ambarwati Ambarwati, Ica Narayani Pramudaningsih</i>	PDF 253-258



### ACCREDITATION



### JOURNAL TEMPLATE



### COUNTER

 22,197	 33
 696	 29
 56	 29
 49	 23
 39	 22

Pageviews: 61,809



### JOURNAL HELP

### USER

<b>Influence Akupresur At The LI 4 And Sp 6 To A Decrease In Pain Dysminorrhea</b>  Ambarwati Ambarwati, Ica Narayani Pramudaningsih	PDF 253-258
<b>The Correlation Number Of Parity On Third Trimester Of Pregnancy To Anxiety In Confronting Birth Labour</b>  Susanti Susanti, Ika Novita Sari	PDF 259-264
<b>The Effectiveness of Warm Water Soak Therapy on Feet and Classical Music Therapy on Blood Pressure of Hypertensive Clients at Puskesmas Ijen Bondowoso</b>  Handono Fatkhur Rahman, Saipul Bahri, Heri Siswanto	PDF 265-274
<b>The Relationship Between Postpartum Depression and Breastfeeding</b>  Uliyatul Laili, Rizki Amalia	PDF 275-279
<b>Patients' Perceptions of Compliance with Tuberculosis Medication in Lamongan</b>  Devi Ristian Octavia, Primanitha Ria Utami	PDF 280-286
<b>The Level Of Knowledge on the Use Of NSAIDs As Analgesic For Dysmenorrhea Case In Faculty of Health Universitas Muhammadiyah Lamongan</b>  Primanitha Ria Utami, Devi Ristian Octavia, Selly Septi Fandinata	PDF 287-294
<b>Lift The Flap Storybook to Grow and Raise Awareness about Sugar Intake in Early Childhood and Their Parents</b>  Diana Dwi Jayanti, Zainal Muttaqin, Dinar Mahdalena Leksana	PDF 295-303
<b>The Relationship Of Mobilization To Episiotomy Wound Healing In Post Partum Mothers At Palembang Ar-Rasyid Hospital</b>  Meta Rosdiana, Putinah Putinah	PDF 304-310
<b>Validation Of The Cadmium Analysis Method (Cd) by Atomic Absorption Spectrophotometry in Traditional Scrub</b>  Eddy Suhardiana, Srie Rezeki Nur Endah	PDF 311-318

### USER

Username   
Password   
 Remember me

### LANGUAGE

Select Language  
English

### JOURNAL CONTENT

Search   
Search Scope  
All

- Browse
- » By Issue
  - » By Author
  - » By Title
  - » Other Journals

### CURRENT ISSUE

ATOM 1.0  
RSS 2.0  
RSS 1.0

### FONT SIZE



# JURNAL MIDPRO

ISSN : 2086-2792 (Print) ISSN : 2684-6764 (Online)

Sekolah Vokasi Program Studi Diploma III Kebidanan Universitas Islam Lamongan



- HOME
- ABOUT
- LOGIN
- REGISTER
- SEARCH
- CURRENT
- ARCHIVES
- INDEXING
- ANNOUNCEMENT
- JOURNAL HISTORY

Home > About the Journal > Editorial Team

## Editorial Team

### EDITOR IN CHIEF

Husnul Husnul Muthoharoh, (Google Scholar) Vocational Program of Midwifery, Universitas Islam Lamongan, Indonesia

### EDITORIAL TEAM

- Widiharti Widiharti, (Google Scholar) Universitas Muhammadiyah Gresik
- Nur Aini, (Google Scholar) The Faculty of Health, Universitas Muhammadiyah Malang
- Rizki Amalia, (Google Scholar) Universitas Nahdlatul Ulama Surabaya
- Yulizawati Yulizawati, (Google Scholar) Universitas Andalas Padang

### MANAGING EDITOR

Fitriana Ikhtiarinawati Fajrin, (Google Scholar) Vocational Program of Midwifery, Universitas Islam Lamongan, Indonesia

- OPEN JOURNAL SYSTEMS
- Editorial Board
- Reviewers
- Peer Review Process
- Section Policies
- Focus and Scope
- Author Guidelines
- Publication Ethics
- Online Submission
- Open Access Policy
- Author(s) fee
- Copyright Notice
- Login
- Contact





# JURNAL MIDPRO

ISSN : 2086-2792 (Print) ISSN : 2684-6764 (Online)

Sekolah Vokasi Program Studi Diploma III Kebidanan Universitas Islam Lamongan



- HOME
- ABOUT
- LOGIN
- REGISTER
- SEARCH
- CURRENT
- ARCHIVES
- INDEXING
- ANNOUNCEMENT
- JOURNAL HISTORY

Home > About the Journal > People

## People

### REVIEWER

Dito Anurogo, (SINTA ID: 6198330) Faculty of Medicine and Health Science, Muhammadiyah University of Makassar

Abdul Muhith, (SCOPUS ID: 57202905408), STIKES Majapahit, Indonesia

Nur Aini, (Google Scholar) The Faculty of Health, Universitas Muhammadiyah Malang

Dewi Ariani, (SINTA ID: 5983306) Universitas Brawijaya Malang, Indonesia

Sunarti sunarti, (SCOPUS ID:57211045283), Universitas Ahmad Dahlan, Yogyakarta, Indonesia

Erli Zainal, (SCHOLAR ID) Akademi Kesehatan Yayasan Saptta Bakti, Bengkulu, Indonesia

Indah Mauludiyah, (SINTA ID: 6104393) STIKES Kendedes Malang, Indonesia

Faridah Faridah, (SINTA ID:6702915) STIKES Hang Tuah Surabaya, Indonesia

Rizka Ayu Setyani, (SINTA ID: 6103670) Universitas Respati Yogyakarta, Indonesia

Ilin Nilawati, (SINTA ID: 6115079) Akademi Kesehatan Yayasan Saptta Bakti, Bengkulu, Indonesia

Kholifatul Ummah, (Scholar ID) Universitas Dr. Soetomo Surabaya, Indonesia

### OPEN JOURNAL SYSTEMS

[Editorial Board](#)

[Reviewers](#)

[Peer Review Process](#)

[Section Policies](#)

[Focus and Scope](#)

[Author Guidelines](#)

[Publication Ethics](#)

[Online Submission](#)

[Open Access Policy](#)

[Author\(s\) fee](#)

[Copyright Notice](#)

[Login](#)

[Contact](#)



### ACCREDITATION

SERTIFIKAT



## Phytochemical Screening And Tyrosinase Inhibition Activity Of Leaves *Cassia siamea* L.

<sup>K</sup>Munawarohthus Sholikha<sup>1</sup>, Ainun Wulandari<sup>2</sup>

<sup>1,2</sup>Departemen Farmasi, Fakultas Farmasi, Institut Sains dan Teknologi Nasional (ISTN)

Email Korespondensi (<sup>K</sup>): [mona.farmasi@istn.ac.id](mailto:mona.farmasi@istn.ac.id)<sup>1</sup>

### ABSTRACT

*Cassia siamea* L. has been used traditionally as medicine because it has chemical constituents such as saponins, anthraquinones, alkaloids, flavonoids, tannins, terpenoids and steroids. Indonesia, which is located in a tropical region with high temperatures and ultraviolet radiation, can cause skin disorders such as hyperpigmentation due to excess melanin synthesis. Tyrosinase enzymes can prevent or inhibit melanin formation. The purpose of this study was to determine the chemical content of *Cassia siamea* L. leaves extract with various solvents and the inhibitory activity of tyrosinase enzymes. In this study, the leaves of *Cassia siamea* L. were macerated with methanol as a solvent. Dry methanol extract was fractionated using the liquid-liquid method using aquadest, butanol and chloroform as solvents. The four dried extracts were then phytochemical screening to determine their chemical content. Tyrosinase inhibition test was carried out in vitro with L-Dopa as a substrate using ELISA plate well reader at concentrations of 100, 1000 and 10000 ppm with three repetitions (triplo). Methanol extract had percent inhibition of  $19.993 \pm 1.125\%$ ,  $28.984 \pm 0.624\%$ ,  $57.164 \pm 0.623\%$ , respectively; butanol fraction  $25.914 \pm 0.541\%$ ,  $32.566 \pm 0.767\%$ ,  $52.120 \pm 1.616\%$ ; chloroform fraction  $19.920 \pm 0.730\%$ ,  $26.425 \pm 0.937\%$ ,  $62.865 \pm 0.167\%$  and aquadest fraction  $23.830 \pm 0.879\%$ ,  $30.885 \pm 0.778\%$ ,  $67.471 \pm 0.352\%$ . Kojic acid had tyrosinase activity and acts more active than the other extract and fraction with  $91.155 \pm 0.228\%$  at 500 ppm. The aquadest fraction had the highest percent inhibition ( $67.471 \pm 0.352\%$ ) at 10000 ppm which was thought to be due to the presence of alkaloids, tannins, flavonoids, steroids and terpenoids from the results of the phytochemical screening test.

Keywords : *Cassia siamea* L; phytochemical screening; tyrosinase

### Article history :

Received: 17 Juni 2020

Received in revised form: 15 September 2020

Accepted: 10 October 2020

Available online: 1 December 2020



licensed by [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).

## INTRODUCTION

Indonesia is a country located in a tropical region which is famous for its high temperatures and ultraviolet (UV) radiation at the highest level. Exposure to UV rays for a long time with frequent frequency can cause skin irritation such as darkening of the skin color. The brown color of the skin is the result of excessive melanin formation<sup>1</sup>. Hyperpigmentation is the most common pigment disorder of the skin. An increase in melanin synthesis or an uneven distribution of melanin can cause hyperpigmentation or patches of skin. The process of forming melanin compounds (melanogenesis) occurs with the help of biocatalysts, especially the enzyme tyrosinase<sup>2</sup>.

In the melanogenesis process, tyrosinase acts as a catalyst in two different reactions, namely the hydroxylation of tyrosine to dihydroxy-phenylalanine (L-DOPA) and the oxidation of L-DOPA to dopakuinone. Dopakuinone compounds have very high reactivity so they can polymerize spontaneously to form dopaque which then becomes melanin<sup>3</sup>. Tyrosinase in skin tissue is activated by solar UV radiation, thereby accelerating the process of melanin production. Tyrosinase enzymes can prevent or inhibit melanin formation<sup>1</sup>.

One way to lighten skin color is by inhibiting melanin formation through inhibition of the tyrosinase enzyme. Bleach acts as an inhibitor of melanin production and is known as a competitive tyrosinase inhibitor. Various tyrosinase inhibitors are found in cosmetic ingredients including hyaluronic, arbutin, kojic acid, mercury, and hydroquinone. This compound has a very large bleaching power, but is dangerous because it is carcinogenic<sup>4</sup>. The discovery of natural ingredients that are safe for human health, one of which is looking for tyrosinase inhibitors found in nature.

*Cassia siamea* L. has been used traditionally as medicine and in previous studies had an IC<sub>50</sub> antioxidant activity of 144.12 µg/mL<sup>5</sup>. *Cassia siamea* L. potential as a medicinal substance is thought to have chemical constituents such as saponins, anthraquinones, alkaloids, flavonoids, tannins, terpenoids and steroids<sup>6</sup>. The total phenolic content (TPC) that has been carried out in previous studies was 88.5 mgGAE/g, while the total flavonoids content (TPC) obtained were 35.6 mgRE/g<sup>7</sup>. Flavonoids with their antioxidant effects play a role in inhibiting the tyrosinase enzyme. The purpose of this study was to determine the chemical content of *Cassia siamea* L. leaves extract with various solvents and the inhibitory activity of tyrosinase enzymes.

## METHODS

*Cassia siamea* L. dry leaves was collected from Balai Penelitian Tanaman Rempah dan Obat (Balitro) which has been determined at the Botanical Garden Plant Conservation Center, Lembaga Ilmu Pengetahuan Indonesia (LIPI). Chemical reagents such as methanol 75%, L-DOPA (Sigma), tyrosinase enzyme (Sigma), kojic acid (Sigma), chloroform, butanol, aquadest, HCl 2 N, Dragendorff reagent, Mayer reagent, Wagner reagent, Bouchardat reagent, HCl p, NaNO<sub>2</sub> 5%, AlCl<sub>3</sub> 10%, NaOH 1N, FeCl<sub>3</sub> 1%, sodium hydroxide NaOH 2N, ether, H<sub>2</sub>SO<sub>4</sub>, potassium dihydrogen phosphate, dimethyl sulfoxide

(DMSO) (Merck), phosphate buffer (pH 6.5). ELISA plate well reader was used for tyrosinase inhibition assay.

### **Sample Preparation, Extraction, and Fractionation**

*Cassia siamea* L. dry leaves was ground to obtain 500 gram sample powder to the extraction process. Methanol 75% 5 L was used as the solvent in the maceration extraction of the samples for 3x24 hours. The crude methanol extracts were then dried using a rotary evaporator. Liquid-liquid fractionation was conducted using aquadest, buthanol, chloform to the methanol extract to obtain fractions with different polarities then dried using a rotary evaporator. The four dried extracts were then phytochemical screening to determine their chemical content and tyrosinase inhibition assay.

### **Phytochemical Screening**

#### **Alkaloids**

0.25 grams of sample was added with 0.5 mL of 2N hydrochloric acid and 4.5 mL of distilled water, heated over a water bath for 2 minutes, cooled and then filtered. The filtrate as much as 3 drops was transferred into three test tubes. To tube 1 are added 2 drops of Wagner solution. If there is a brown to black sediment, the powder contains alkaloids. Tube 2 is added with 2 drops of Mayer reagent. If a white precipitate is formed, the powder contains alkaloids. Tube 3 is added 2 drops of Dragendorff reagent. If a brown sediment is formed, the attack contains alkaloids.

#### **Flavonoids**

0.5 gram of sample was extracted with 50 mL of hot water, then filtered with filter paper until the filtrate was obtained. 2.5 mL of the filtrate is poured into the test tube. The filtrate was added with 0.5 mL of 5% NaNO<sub>2</sub> solution and 0.5 mL of 10% AlCl<sub>3</sub> and then shaken. 1 mL of 1N NaOH was added slowly into the filtrate through the test tube wall. The presence of flavonoids indicates a red or orange color

#### **Saponins**

0.5 gram of sample was extracted with 100 mL of hot water, then filtered with filter paper until the filtrate was obtained. 5 mL of the filtrate is poured into a test tube then shaken vigorously for 10 seconds. If a foam is formed that is stable for not less than 10 minutes and does not disappear with the addition of 1 drop of 2 N hydrochloric acid, it indicates the presence of saponins

#### **Tannins**

0.5 gram of sample was extracted with 50 mL of hot water, then filtered with filter paper until the filtrate was obtained. 2.5 mL of the filtrate is poured into the test tube. Then add 2-3 drops of 1% iron (III) chloride solution into the test tube containing the filtrate. The presence of the phenol group is indicated by the formation of a green-blue-black color

#### **Steroids & Terpenoids**

0.5 gram of sample was macerated using 10 mL ether for 2 hours then filtered with filter paper. 2.5 mL of the filtrate is poured into the test tube. 0.5 mL H<sub>2</sub>SO<sub>4</sub> is poured into a test tube containing the



filtrate. Then drop the liebermann-burchard reagent slowly through the wall. The formation of green violet or blue color indicates the presence of steroids / triterpenoids

### Tyrosinase Inhibitory Assay

Tyrosinase inhibitory activity was evaluated based on inhibition of the sample (dilute DMSO) to diphenolase activity. The assay was carried out using ELISA plate well reader with tyrosinase enzyme, L-DOPA as the substrates, phosfat buffer pH 6.5, with three repetitions (triplo)<sup>17</sup>. Kojic acid was used as a positive control. The percentage of tyrosinase inhibitory activity can be calculated by the following formula:

$$\text{Inhibition (\%)} = \left[ 1 - \frac{A-C}{B-D} \right] \times 100\%$$

Where,

A: Absorbance of the sample

B: Absorbance of blank

C: Absorbance of sampel control

D: Absorbance of blank control

## RESULTS

The extraction process using methanol as a solvent is based on previous research that the leaves of *Cassia siamea* L. had the lowest IC<sub>50</sub> value in the antioxidant activity test<sup>5</sup>. The low IC<sub>50</sub> value of antioxidants shows the effectiveness in counteracting free radicals from UV rays which indirectly affect the inhibition of the tyrosinase enzyme. The maceration method is chosen because it has advantages, namely in the fast processing process, the method of working and the equipment used is simple, relatively easy and inexpensive<sup>8</sup>. Below is the yield of the extract and fraction of the leaves of *Cassia siamea* L.

Table 1. the yield of the extract and fraction of leaves of *Cassia siamea* L.

Simplicia	Fraction	Powder Weight (g)	Extract Weight (g)	Yield %
<i>Cassia siamea</i> L. Leaves	Methanol	500	70.5	14.1
	Chloroform	40	9.1	22.75
	Aquadest	40	12.2	30.5
	Butanol	40	5	12.5

Phytochemical screening was carried out using the previous method<sup>9</sup>. The most important types of phytochemicals found in extract and fraction of leaves of *Cassia siamea* L. are alkaloids, flavonoids, tannins, steroids, terpenoids positively detected during phytochemical confirmation as shown in Table 2.

Table 2. Phytochemical screening test results of leaves of *Cassia siamea* L.

Sample	Phytochemical Tested	Test Result	
Methanol Extract	Saponin	-	
	Alkaloids	Mayer	-
		Wagner	-
		Dragendorff	+
	Tannins	-	
	Flavonoids	+	
Steroids & terpenoids	-		
Butanol Fraction	Saponin	-	
	Alkaloids	Mayer	+
		Wagner	-
		Dragendorff	+
	Tannins	+	
	Flavonoids	+	
Steroids & terpenoids	+		
Chloroform Fraction	Saponin	-	
	Alkaloids	Mayer	-
		Wagner	-
		Dragendorff	-
	Tannins	-	
	Flavonoids	+	
Steroids & terpenoids	-		
Aquadest Fraction	Saponin	-	
	Alkaloids	Mayer	+
		Wagner	-
		Dragendorff	+
	Tannins	+	
	Flavonoids	+	
Steroids & terpenoids	+		

Tyrosinase inhibitory activity was evaluated to measure the ability of samples as lightening agents. Tyrosinase inhibition is correlated with the decrease of melanogenesis on the skin since this enzyme is responsible for hyperpigmentation in humans<sup>10</sup>. In this study, using the L-DOPA substrate because this substrate with the tyrosinase enzyme will produce dopaque which can then form melanin to prevent it by tyrosinase inhibitors. The formation of dopaid products is characterized by the appearance of a brown color which can cause inhibition of the substrate-tyrosinase reaction, resulting in reduced dopaque production which is indicated by a decrease in the intensity of the brown color. This dopaque formation uses a microplate reader using 96 wells microplates at a wavelength of 492 nm<sup>11</sup>.

Table 3. Tyrosinase inhibitory of leaves of *Cassia siamea* L.

Sample	Concentration (ppm)	Tyrosinase Inhibition (%)
Methanol Extract	100	19.993 ± 1.125
	1000	28.984 ± 0.624
	10000	57.164 ± 0.623
Butanol Fraction	100	25.914 ± 0.541
	1000	32.566 ± 0.767
	10000	52.120 ± 1.616
Chloroform Fraction	100	19.920 ± 0.730
	1000	26.425 ± 0.937
	10000	62.865 ± 0.167
Aquadest Fraction	100	23.830 ± 0.879
	1000	30.885 ± 0.778
	10000	67.471 ± 0.352
Kojic Acid	500	91.155±0.228

## DISCUSSION

Each phytochemical showed potency towards some biological action; for example, flavonoids with their antioxidant effects play a role in inhibiting the tyrosinase enzyme<sup>12</sup>. Tannins also have the ability to act as anti-tyrosinase because they inhibit the biosynthesis process of melanin so that an increase in melanin production does not occur after exposure to UVB rays<sup>13</sup>. Inhibitors of other classes, such as terpenes, steroids, alkaloids, have also been reported in the tyrosinase inhibitory activity<sup>14</sup>.

Investigation of tyrosinase inhibitory activity showed that aquadest fraction had the highest percent inhibition ( $67.471 \pm 0.352\%$ ) at 10000 ppm compared to the other extract and fraction. The presence of alkaloids, tannins, flavonoids, steroids and terpenoids from the results of the phytochemical screening test of aquadest fraction play a role in inhibiting the tyrosinase enzyme. Kojic acid has tyrosinase activity and acts more active than the other extract and fraction with  $91.155 \pm 0.228\%$  at 500 ppm. This was affected by the extract conditions, which was a crude extract and fraction which had not been purified, so it was suspected there were other compounds that did not act as tyrosinase inhibitors. Kojic acid has an inhibitory effect of tyrosinase enzyme and the greatest stability in cosmetic products<sup>15</sup>, but it is carcinogenic and its use in high concentration can damage the skin<sup>16</sup>.

## CONCLUSIONS

The highest yield of aquadest fraction amounted to 30.5%. Concentration 100, 1000, 10000 ppm shows that methanol extract had percent inhibition of  $19.993 \pm 1.125\%$ ,  $28.984 \pm 0.624\%$ ,  $57.164 \pm 0.623\%$ , respectively; butanol fraction  $25.914 \pm 0.541\%$ ,  $32.566 \pm 0.767\%$ ,  $52.120 \pm 1.616\%$ ; chloroform fraction  $19.920 \pm 0.730\%$ ,  $26.425 \pm 0.937\%$ ,  $62.865 \pm 0.167\%$  and aquadest fraction  $23.830 \pm 0.879\%$ ,  $30.885 \pm 0.778\%$ ,  $67.471 \pm 0.352\%$ . Kojic acid had tyrosinase activity and acts more active than the other extract and fraction with  $91.155 \pm 0.228\%$  at 500 ppm. The aquadest fraction had the highest percent inhibition ( $67.471 \pm 0.352\%$ ) at 10000 ppm which was thought to be due to the presence of alkaloids, tannins, flavonoids, steroids and terpenoids from the results of the phytochemical screening test.

Aquadeest fraction of leaves *Cassia siamea* L. can be used as active skin lightening ingredients and formulated into cosmetic because have highest yield and inhibitory effect of tyrosinase enzyme.

### ACKNOWLEDGMENTS

The authors are thankful to acknowledge the Ministry of Research, Technology, and Higher Education, Republic of Indonesia, for support via Research Grant (26/E1/KPT/2020).

### REFERENCES

1. Sagala Z, Pratiwi RW, Azmi NU, Maap. Uji Aktivitas Inhibisi terhadap Enzim Tirosinase dari Ekstrak Etanol Daun Pepaya ( *Carica papaya* L .) Secara In Vitro. J Penelit Farm Indones. 2019;7(2):34–8.
2. Rauf, Afrisusnawati., Ningsi, Surya., Nurdin, Hasmawati., 2017. Potensi Penghambatan Tirosinase Ekstrak Etanol Daun Tomat (*Lycopersicon esculentum* Mill, var. Pyriforme Alef). J Farmasi Uin Alauddin.2019;7(1)
3. Juwita NK, Djajadisastra J. Uji Penghambatan Tirosinase Dan Yang Mengandung Ekstrak Kulit Batang Nangka (*Artocarpus heterophyllus*). 2011;8(3):127–40.
4. Noor SU, Faridah, Magdalena P. Uji Aktivitas Inhibisi Enzim Tirosinase In-Vitro Krim Ekstrak Akar Manis (*Glycyrrhiza glabra* L.). J Ilmu Kefarmasian Indones. 2018;16(2):150–8.
5. Mehta JP, Parmar PH, Vadia SH, Patel MK, Tripathi CB. In-vitro antioxidant and in-vivo anti-inflammatory activities of aerial parts of *Cassia* species. Arab J Chem [Internet]. 2017;10:S1654–62. Available from: <http://dx.doi.org/10.1016/j.arabjc.2013.06.010>
6. Kamagaté M, Koffi C. Ethnobotany, phytochemistry, pharmacology and toxicology profiles of *Cassia siamea* Lam. J Phytopharm [Internet]. 2014;3(1):57–76. Available from: [http://www.phytopharmajournal.com/vol3\\_issue1\\_09.pdf](http://www.phytopharmajournal.com/vol3_issue1_09.pdf)
7. Kaisoon O, Siriamornpun S, Weerapreeyakul N, Meeso N. Phenolic compounds and antioxidant activities of edible flowers from Thailand. J Funct Foods [Internet]. 2011;3(2):88–99. Available from: <http://dx.doi.org/10.1016/j.jff.2011.03.002>
8. Kurniasari A, Anwar E, Djajadisastra J. Potensi Ekstrak Biji Coklat (*Theobroma cacao* Linn) sebagai Inhibitor Tirosinase untuk Produk Pencerah Kulit. J Kefarmasian Indones. 2018;8(1):34–43.
9. Manurung H, Aryani R, Nugroho RA, Sari YP, Chernovita R, Auliana. Phytochemical analysis and antioxidant activity of leaves extracts of endemic plant Jahe Balikpapan (*Etlingera balikpapanensis* A.D. Poulsen). Int J Sci Technol Res. 2019;8(9):308–13.
10. Chang TS. An updated review of tyrosinase inhibitors. Int J Mol Sci. 2009;10(6):2440–75.
11. Sari RK, Utami R, Batubara I, Carolina A, Febriany S. Aktivitas Antioksidan dan Inhibitor Tirosinase Ekstrak Metanol Mangium (*Acacia mangium*) (Antioxidant and Tyrosinase Inhibitor Activities of Methanol Extracts of *Acacia mangium*). J Ilmu Teknol Kayu Trop. 2015;13(1).
12. Ohguchi K, Tanaka T, Kido T, Baba K, Inuma M, Matsumoto K, et al. Effects of hydroxystilbene derivatives on tyrosinase activity. Biochem Biophys Res Commun. 2003;307(4):861–3.
13. Siahaan ER, Pangkahila W, Wiraguna AAGP. Krim ekstrak kulit delima merah (*Punica*

- granatum) menghambat peningkatan jumlah melanin sama efektifnya dengan krim hidrokuinon pada kulit marmut (*Cavia porcellus*) betina yang dipapar sinar UVB. *J Biomedik*. 2017;9(1):7–13.
14. Chiou SY, Ha CL, Wu PS, Yeh CL, Su YS, Li MP, et al. Antioxidant, Anti-tyrosinase and Anti-inflammatory activities of oil production residues from *Camellia tenuiflora*. *Int J Mol Sci*. 2015;16(12):29522–41.
  15. Dolorosa MT, Nurjanah, Purwaningsih S, Anwar E, Hidayat T. Tyrosinase inhibitory activity of *Sargassum plagyophyllum* and *Eucheuma cottonii* methanol extracts. *IOP Conf Ser Earth Environ Sci*. 2019;278(1).
  16. Miyazawa M, Tamura N. Inhibitory compound of tyrosinase activity from the sprout of *Polygonum hydropiper* L. (Benitade). *Biol Pharm Bull*. 2007;30(3):595–7.
  17. Batubara I, Darusman LK, Mitsunaga T, Rahminiwati M, Djauhari E. Potency of Indonesian medicinal plants as tyrosinase inhibitor and antioxidant agent. Vol. 10, *Journal of Biological Sciences*. 2010. p. 138–44.